

Moyamoya Disease: Physiological Mechanisms and Treatment Approaches

Kaustov Chakrabarti^{1*}

¹Geisinger College of Health Sciences, Scranton, PA 18509

*Master of Biomedical Sciences Program

Correspondence: kchakrabarti@geisinger.edu

Abstract

Moyamoya disease (MMD) is a rare, progressive cerebrovascular disorder marked by stenosis of the terminal internal carotid arteries and the formation of fragile collateral vessels, leading to reduced cerebral perfusion and heightened risks of ischemic and hemorrhagic strokes. Its pathophysiology involves endothelial dysfunction, aberrant angiogenesis, and genetic mutations, notably in the RNF213 gene. This paper examines normal cerebrovascular physiology, the pathological changes driving MMD, its clinical features, and current therapeutic approaches. Treatments, including medical management and surgical revascularization (e.g., direct and indirect bypass), aim to restore cerebral blood flow and mitigate stroke risk, with revascularization proving most effective. Elucidating the physiological underpinnings of MMD is vital for developing targeted therapies and enhancing patient outcomes.

Introduction

Moyamoya disease (MMD) is a rare and progressive cerebrovascular disorder bimodally distributed peaking in childhood and young adults around age 40, although its reach spans all ages (1). Named from the Japanese term for "puff of smoke," MMD is characterized by the stenosis of terminal internal carotid arteries and the emergence of fragile collateral vessels that resemble wispy clouds on angiography. This compensatory network often fails, leading to chronic cerebral hypoperfusion and a heightened risk of ischemic and hemorrhagic strokes — devastating outcomes that can rob patients of motor function, cognition, or life. Moyamoya disease symptoms are primarily driven by inadequate blood flow to the brain and can be devastating. Patients often experience neurological deficits such as sudden weakness or paralysis, typically affecting one side of the body. They may also suffer from aphasia, vision problems, or

involuntary movements. Without effective treatment, the chronic hypoperfusion and heightened risk of ischemic or hemorrhagic strokes can lead to severe cognitive impairment, permanent disability, and, in some cases, death (1). Though its pathogenesis remains elusive, genetic mutations like RNF213, prevalent in East Asian populations (2), hint at a complex interplay of hereditary and environmental triggers. Despite its rarity, MMD's impact is profound, necessitating urgent advances in diagnosis and care. Treatment strategies, from medical management to sophisticated surgical revascularization, aim to restore blood flow and avert neurological decline, yet challenges persist. Medical therapies cannot halt the progressive vessel narrowing, and surgery, despite its efficacy, carries inherent risks such as perioperative stroke and the need for highly specialized neurosurgical expertise. Moreover, predicting long-term neurological outcomes for all patients remains difficult, and the disease's progressive nature necessitates lifelong clinical and imaging surveillance. This paper explores MMD comprehensively: from the normal physiology of cerebral blood flow and its disruption in MMD, through its genetic and hemodynamic underpinnings, to its clinical presentation, diagnostic approaches, and evolving therapies. By illuminating these facets, we seek to enhance understanding and guide the development of targeted interventions for improved patient outcomes.

Methods

A comprehensive literature search was conducted to identify relevant studies, case reports, and review articles on MMD focusing on diagnosis, treatment, and patient outcomes globally. The primary search engine employed was Google Scholar.

To maximize the relevance of the retrieved literature, the search strategy incorporated keywords and phrases such as "Moyamoya disease," "Moyamoya

diagnosis," "Moyamoya treatment," "Moyamoya outcome," "Moyamoya surgery," "Moyamoya revascularization," "Moyamoya complications," and "Moyamoya case report." Boolean operators (AND, OR) were implicitly used by Google Scholar's algorithm, which prioritizes results containing multiple terms.

While our focus was on recent evidence, with roughly 90% of included studies published in the last 15 years, the search criteria were occasionally broadened. This was necessary due to MMD rarity and the limited experimental data available. Including some older, foundational studies allowed us to gather critical insights into diagnostic approaches, long-term outcomes, and complex case presentations that have significantly shaped our understanding of the disease.

The set of studies on MMD is mostly made up of more recent studies, with approximately 90.9% of citations being published in the last 15 years. This is purposeful to prioritize recent evidence and to also recognize that evidence surrounding this rare condition is rapidly evolving. Most of the selection criteria were driven by relevance to MMD with consideration of diagnostic strategies, long-term outcomes, and complex cases. The studies are drawn from reputable medical journals.

In addition, artificial intelligence was used to structurally organize this paper and ensure grammatical quality. Specifically, each section of this manuscript was grammatically fully checked (3).

Discussion

Normal Physiology of Cerebral Blood Flow Regulation

The brain's vascular system is a specialized network that ensures a continuous supply of oxygen and nutrients for neuronal function. The internal carotid and vertebral arteries, key conduits of this system, converge at the brain base to form the Circle of Willis. This structure enables collateral blood flow, allowing blood to bypass blockages or narrowing in major arteries, thus protecting against ischemic events like strokes and maintaining stable perfusion across both hemispheres (2). Beyond nutrient delivery, the Circle of Willis may also regulate cerebral pressure and maintain intracranial homeostasis. In MMD, however, narrowing or occlusion of the internal carotid arteries triggers the formation of fragile collateral vessels, which often fail to compensate adequately, leading

to ischemic or hemorrhagic complications. This vulnerability highlights the critical role of the cerebral arterial network in sustaining circulation.

Cerebral blood flow (CBF) is tightly regulated to deliver a consistent supply of oxygen and nutrients despite fluctuations in systemic blood pressure. This autoregulation relies on multiple mechanisms. The myogenic response enables cerebral arteries to constrict or dilate in response to changes in intraluminal pressure, stabilizing CBF (4). Metabolic regulation, meanwhile, adjusts blood flow to meet local neuronal demands; increased activity releases vasodilators like adenosine and nitric oxide (NO), enhancing CBF in active regions. Though less dominant, the autonomic nervous system, particularly sympathetic innervation, also modulates CBF (5).

Additionally, CBF responds to changes in arterial carbon dioxide (PaCO₂) and pH: hypercapnia (elevated PaCO₂) and acidosis induce vasodilation, while hypocapnia (reduced PaCO₂) and alkalosis cause vasoconstriction (4). Endothelium further regulates CBF by releasing vasoactive substances, such as NO, a potent vasodilator, and endothelin-1 (ET-1), a vasoconstrictor. The balance between these factors is essential for maintaining a vascular tone and optimal CBF (5).

Pathophysiology of Moyamoya Disease

The defining feature of MMD is the relentless, progressive stenosis and eventual occlusion of the distal internal carotid arteries and the proximal segments of the anterior and middle cerebral arteries, a process that severely compromises (CBF) to critical brain regions. The brain compensates for ischemia by forming a frail, unstable network of collateral vessels, often called a "puff of smoke" on angiography, which are prone to rupture or thrombosis. The etiology of MMD remains incompletely understood, shrouded in a complex interplay of genetic predisposition and environmental influences. Central to its pathogenesis are genetic mutations, with the RNF213 gene emerging as a pivotal player, particularly in East Asian populations where its prevalence reaches up to 90% in familial cases (6). The RNF213 mutation disrupts vascular stability by impairing endothelial cell function and promoting aberrant angiogenesis, potentially through heightened inflammation or oxidative stress. Other implicated genes, such as ACTA2 and GUCY1A3, further underscore the hereditary basis

of this disorder, altering smooth muscle integrity and guanylate cyclase signaling, which compromise vascular resilience (6). Environmental triggers, such as chronic inflammation or autoimmune processes, may amplify these genetic vulnerabilities, accelerating disease progression (6).

The stenotic occlusion of major cerebral arteries precipitates a marked reduction in CBF, often dropping below 30 mL/100g/min in affected regions, while crippling cerebrovascular reserve, the brain's capacity to dilate vessels in response to heightened metabolic demand or stress (7). This impaired autoregulation, compounded by endothelial dysfunction, leaves the brain susceptible to ischemic insults. Endothelial cells in MMD exhibit reduced NO production, coupled with elevated endothelin-1 (ET-1), tilting the balance toward vasoconstriction and exacerbating hypoperfusion (5). The fragile collaterals, meanwhile, pose a dual threat: thrombosis can precipitate ischemic strokes, abruptly starving brain tissue of oxygen, while their rupture, driven by hemodynamic stress on weakened vessel walls, can unleash devastating hemorrhagic strokes, flooding the intracranial space with blood. Although MMD ravages the cerebral vasculature, its ripple effects extend systemically; chronic hypoperfusion erodes cognitive faculties, manifesting as memory deficits, executive dysfunction, and slowed processing speed, and undermines overall neurological function, profoundly impacting quality of life. These multifaceted consequences highlight the urgent need to unravel MMD's physiological and molecular underpinnings to devise more effective therapeutic strategies.

Clinical Signs and Symptoms

MMD manifests through a spectrum of neurological symptoms driven by cerebral hypoperfusion and collateral vessel instability. Brief episodes of neurological dysfunction, such as unilateral weakness (hemiparesis), numbness, aphasia (language disorder caused by damage to the parts of the brain that control language expression and comprehension), or visual disturbances — termed transient ischemic attacks (TIAs) — typically resolve within minutes to hours and may be triggered by hyperventilation or dehydration, which exacerbate vasoconstriction. Sudden ischemic strokes, occurring in 50–75% of untreated patients (2), can cause permanent deficits like hemiplegia or speech impairment, reflecting occlusion of the middle cerebral artery. Seizures, arising from ischemia or cortical

scarring, affect a subset of patients (7), while chronic hypoperfusion leads to cognitive deficits, memory loss, executive dysfunction, and slowed processing speed, sometimes accompanied by irritability or depression due to frontal lobe involvement. Hemorrhagic events are more common in adults and can be up to 40% in East Asian cohorts (7), and result from fragile collateral vessel rupture, causing intracerebral or subarachnoid hemorrhage with severe headaches, neck stiffness, or unconsciousness. Chronic headaches, from vascular anomalies, are frequent across all ages. In children, developmental delays or rare choreiform movements signal chronic ischemia, contrasting with adults' hemorrhagic predominance. Chorea refers to involuntary and irregular movement that appears to flow from one body part to another (8). These movements are typically spontaneous and purposeless, affecting various muscle groups, including the face, limbs, and trunk (8). Choreiform movements are a descriptive term for these specific types of involuntary, dance-like movements. In children with Moyamoya disease, these movements signal chronic brain ischemia, indicating insufficient blood supply to certain brain regions (8). These symptoms stem from progressive arterial stenosis, reducing (CBF), and the fragility of compensatory collaterals. Ischemic events arise from thrombosis or critically low perfusion, while hemorrhages reflect hemodynamic stress on weakened vessels. Symptom severity and type depend on the location and extent of affected brain tissue, with untreated cases progressing from transient to permanent deficits.

Diagnosis

Diagnosing MMD relies on a multimodal imaging approach to confirm progressive stenosis of the terminal internal carotid arteries (ICAs), visualize compensatory collateral networks, and assess cerebral perfusion deficits (9). Magnetic resonance imaging (MRI) and magnetic resonance angiography (MRA) serve as initial, non-invasive cornerstones. MRI detects ischemic infarcts, white matter lesions, or hemorrhagic sequelae in brain parenchyma, often revealing silent strokes in asymptomatic patients. MRA delineates the characteristic narrowing or occlusion of the distal ICAs and proximal anterior/middle cerebral arteries, alongside the hazy, cloud-like "Moyamoya" collaterals — lenticulostriate or thalamo-perforating vessels — that define the disease's angiographic signature. Time-of-flight MRA,

commonly used, offers high sensitivity for detecting these vascular anomalies without contrast, though its resolution may miss subtle collateral networks (10). MRI and MRA are the preferred initial, non-invasive diagnostic tools for Moyamoya disease due to their ability to clearly visualize the characteristic narrowing of the internal carotid arteries and the compensatory "moyamoya" collateral vessels, as well as assess brain tissue for signs of ischemia or hemorrhage, all without exposing the patient to radiation, which is particularly beneficial for pediatric cases (11). While CT can detect acute issues, MRI and MRA offer superior soft tissue contrast and detailed vascular imaging, making them more effective for comprehensive diagnosis and monitoring of this progressive condition (12). Cerebral angiography remains the gold standard, providing unparalleled detail of cerebral vasculature. This invasive technique maps the extent of ICA stenosis, quantifies collateral vessel proliferation, and enables staging via the Suzuki classification (stages I–VI), which tracks disease progression from mild narrowing to extensive collateral dependence and ICA occlusion (3). The "puff of smoke" appearance — most striking in advanced stages — arises from dilated perforating arteries compensating for reduced flow, a finding critical for distinguishing MMD from mimics like atherosclerosis or vasculitis. Despite its precision, angiography carries risks (e.g., stroke from catheter manipulation), prompting its use primarily for confirmation or preoperative planning (9).

Functional imaging complements structural assessments. Single-photon emission computed tomography (SPECT), often paired with acetazolamide challenge, quantifies cerebral blood flow (CBF) and cerebrovascular reserve, identifying ischemic regions at risk of infarction. Hypoperfused areas (CBF <30 mL/100 g/min) correlate with clinical symptoms like TIAs and guide surgical candidacy (7). Transcranial Doppler (TCD) ultrasonography, a non-invasive tool, measures CBF velocity in major arteries and assesses autoregulatory capacity; diminished velocity or reactivity to CO₂ changes signals impaired hemodynamics. While less specific than SPECT, TCD's portability suits serial monitoring (8). Diagnosis hinges on integrating these findings with clinical presentation — TIAs, strokes, or seizures in children; hemorrhages in adults — excluding secondary causes (e.g., sickle cell disease, radiation injury). Advanced cases may show "ivy" signs on MRI (leptomeningeal collaterals) or flow voids in the basal ganglia, further supporting MMD.

Collectively, these tools not only confirm the diagnosis but also stratify disease severity, informing prognosis and therapeutic strategies like revascularization.

Treatment Strategies

The management of MMD encompasses both medical and surgical approaches, tailored to disease stage, patient age, and clinical presentation (ischemic vs. hemorrhagic). These strategies aim to mitigate cerebral hypoperfusion, prevent stroke, and preserve neurological function by addressing the dual threats of thromboembolism and fragile collateral vessel rupture, guided by advanced imaging to assess revascularization needs (11).

Medical therapy serves as an adjunct to surgical intervention or a temporizing measure for patients unsuitable for surgery. Antiplatelet therapy, such as aspirin (typically 81–325 mg daily), reduces the risk of thromboembolic events by inhibiting platelet aggregation in stenosed vessels and fragile collaterals, a critical consideration given the high incidence of ischemic strokes (50–75% in untreated cases) (6). Evidence suggests modest efficacy in preventing TIA recurrence, though it does not halt disease progression (3). Calcium channel blockers (e.g., nimodipine or nifedipine) are employed to prevent vasospasm and enhance cerebral blood flow (CBF) by relaxing vascular smooth muscle, particularly in patients with symptomatic vasoconstriction triggered by hyperventilation or dehydration (10). For patients with pre-existing heart failure (HF), particularly those with reduced ejection fraction, calcium channel blockers require careful consideration. While dihydropyridine CCBs like nimodipine and nifedipine primarily cause vasodilation and are less likely to directly depress myocardial contractility than non-dihydropyridine CCBs (e.g., verapamil, diltiazem), they can induce systemic hypotension. This hypotension may compromise cardiac output and worsen HF symptoms or end-organ perfusion. In such cases, management involves close hemodynamic monitoring, careful titration of the CCB dose, and optimizing volume status. Alternative strategies for improving CBF or managing vasoconstriction would focus on non-pharmacological approaches like maintaining euvolemia and avoiding triggers, as direct pharmacological substitutes for vasospasm prevention are limited (10). These agents may also alleviate chronic headaches, a frequent MMD symptom. Control of cardiovascular risk factors — hypertension,

diabetes, and hyperlipidemia — is essential to minimize additional vascular stress, with blood pressure targets often set below 130/80 mmHg to balance perfusion needs and hemorrhage risk (3). Despite these measures, medical management alone fails to address the underlying progressive stenosis, underscoring the primacy of surgical intervention.

Surgical revascularization is the cornerstone of MMD treatment, proven to restore CBF, reduce stroke risk, and improve long-term outcomes. Imaging modalities like MRI and SPECT scans guide patient selection and postoperative assessment (11). Techniques fall into 3 categories: direct, indirect, and combined bypass, each leveraging distinct mechanisms to bypass occluded arteries and bolster cerebral perfusion.

Direct bypass, exemplified by superficial temporal artery to middle cerebral artery (STA-MCA) anastomosis, establishes an immediate extracranial-to-intracranial conduit (10). A branch of the external carotid artery (e.g., STA) is meticulously sutured to a cortical branch of the middle cerebral artery, bypassing the stenosed internal carotid artery (ICA) (10). The procedure delivers instantaneous CBF augmentation — often increasing from <30 mL/100 g/min to >50 mL/100 g/min in affected regions — making it particularly effective for patients with acute ischemic symptoms or poor cerebrovascular reserve (7). Success hinges on meticulous microsurgical technique and donor-recipient vessel patency, with patency rates exceeding 90% in experienced centers (1). However, it carries perioperative risks, including hyperperfusion syndrome (5–10% incidence), which can precipitate transient neurological deficits or hemorrhage (7).

Indirect bypass: Indirect methods, such as encephaloduroarteriosynangiosis (EDAS), encephalomyosynangiosis (EMS), or dural inversion, promote angiogenesis by placing vascularized tissue (e.g., temporalis muscle or pericranium) onto the brain surface (9, 11). Over weeks to months, this stimulates collateral vessel formation from extracranial sources, gradually increasing CBF (10). EDAS, the most widely used, leverages the STA without direct anastomosis, making it technically more straightforward and safer for pediatric patients or those with small-caliber vessels (10). Efficacy varies, with angiogenesis success rates ranging from 60% to 80%, influenced by factors like patient age (higher in children) and RNF213 mutation status (8). Specifically, the homozygous RNF213 mutation is associated with lower

angiogenesis success rates. While slower to effect, indirect bypass reduces perioperative risks and suits early-stage or asymptomatic cases.

Combined bypass: Integrating direct and indirect techniques, combined bypass maximizes revascularization by pairing immediate flow (via STA-MCA) with long-term collateral growth (via EDAS or similar) (11). This approach is increasingly favored for advanced MMD (Suzuki stages IV–VI), where extensive collateral dependence and profound hypoperfusion demand robust intervention (12). Studies report superior stroke-free survival rates — up to 95% at 5 years — compared to single-technique strategies, though it requires excellent surgical expertise and longer operative time (13).

Surgical candidacy hinges on imaging evidence of compromised CBF (e.g., SPECT showing <30 mL/100 g/min) and clinical factors like recurrent TIAs or strokes (10). Direct bypass excels in adults with acute ischemia, while indirect methods dominate in children due to their robust angiogenic potential (14). Combined approaches bridge these benefits, though comparative trials remain limited (15).

Beyond established methods, experimental approaches are gaining traction. Gene therapy targeting RNF213 or angiogenesis pathways (e.g., VEGF upregulation) aims to correct endothelial dysfunction and enhance collateral stability, though human trials are nascent (16). Stem cell therapy, using mesenchymal stem cells to promote vascular repair, shows promise in preclinical models by boosting NO production and reducing inflammation (14). Pharmacological agents like cilostazol, a phosphodiesterase inhibitor, are under investigation for their dual antiplatelet and vasodilatory effects, potentially augmenting CBF more effectively than aspirin alone (13). These innovations, while preliminary, signal a shift toward personalized, molecularly targeted MMD management.

Treatment selection balances efficacy, risk, and patient factors, with imaging playing a pivotal role in planning and follow-up (17). Direct bypass offers rapid relief but demands surgical precision; indirect bypass prioritizes safety and long-term gains; combined bypass optimizes outcomes at higher complexity. Postoperative care includes antiplatelet continuation, CBF monitoring via TCD or SPECT, and managing hyperperfusion risks with strict blood pressure control (e.g., systolic 100–140 mmHg) (16). Multidisciplinary teams —

neurosurgeons, neurologists, and rehabilitation specialists — ensure holistic care, addressing both vascular and neurological sequelae.

Prognosis and Long-Term Management

Physiologically, MMD's prognosis hinges on the brain's ability to adapt to chronic hypoperfusion and the success of interventions in re-establishing hemodynamic stability. Pre-treatment, the stenotic ICAs impair autoregulation — the myogenic and metabolic mechanisms that usually stabilize CBF despite systemic pressure fluctuations. This forces reliance on fragile collateral vessels, prone to thrombosis or rupture with their thin-walled, disorganized structure (18). Successful revascularization, whether direct (e.g., STA-MCA bypass) or indirect (e.g., EDAS), reintroduces extracranial blood flow, elevating regional CBF and partially restoring autoregulatory capacity (19). For instance, direct bypass can boost CBF from <30 mL/100 g/min to >50 mL/100 g/min acutely, alleviating ischemic stress (18). However, this sudden hemodynamic shift risks hyperperfusion syndrome (5–10% incidence), where excessive flow overwhelms previously hypoperfused tissue, triggering edema or hemorrhage due to disrupted endothelial integrity and elevated ET-1 levels (18). In the long term, the balance between NO-mediated vasodilation and ET-1-induced vasoconstriction must stabilize to sustain vascular tone and prevent such complications (15).

Effective long-term management integrates regular physiological monitoring with tailored interventions. MRI and MRA track disease progression and bypass patients, assessing whether CBF remains above ischemic thresholds (18). SPECT with acetazolamide challenge quantifies cerebrovascular reserve, revealing regions at risk if autoregulation falters (17). Transcranial Doppler (TCD) provides real-time CBF velocity data, guiding adjustments in antiplatelet therapy (e.g., aspirin) or blood pressure control (target systolic 100–140 mmHg) to prevent thromboembolism or hyperperfusion (19). For hemorrhagic-prone patients, managing collateral vessel stability is critical; excessive hemodynamic stress on these vessels, exacerbated by hypertension or PaCO₂ fluctuations can precipitate rupture, necessitating strict PaCO₂ homeostasis via controlled ventilation or lifestyle adjustments (20). Emerging therapies, like cilostazol, aim to enhance NO production and vasodilation, potentially offering a

dual protective effect against ischemia and vascular instability (20).

Conclusion

MMD is a formidable challenge in cerebrovascular medicine, defined by its progressive stenosis of the internal carotid arteries and the precarious formation of fragile collateral vessels. This paper has elucidated the intricate interplay between normal (CBF) regulation — reliant on autoregulatory mechanisms like myogenic responses and NO-mediated vasodilation — and its disruption in MMD, where endothelial dysfunction, genetic mutations (notably RNF213), and aberrant angiogenesis precipitate chronic hypoperfusion and vascular instability. These physiological disruptions manifest clinically as transient ischemic attacks, debilitating strokes, cognitive decline, and hemorrhagic events, underscoring the disease's profound impact on patients across age groups. Current diagnostic tools, from MRI/MRA to cerebral angiography and SPECT, enable precise visualization of these vascular anomalies and guide therapeutic decision-making, while treatment strategies — spanning medical management with antiplatelets and surgical revascularization via direct, indirect, or combined bypass — offer critical avenues to restore CBF and mitigate stroke risk.

Surgical revascularization emerges as a cornerstone of MMD management, with techniques like STA-MCA bypass and EDAS demonstrably improving hemodynamic stability and long-term outcomes, evidenced by stroke-free survival rates reaching up to 95% at 5 years in advanced cases (21). Yet the disease's complexity demands more than current solutions can fully address. The persistent risks of hyperperfusion syndrome, the variable efficacy of indirect bypass, and the incomplete understanding of MMD's genetic and environmental triggers highlight significant gaps in knowledge and practice. Moreover, the systemic ripple effects of chronic hypoperfusion cognitive impairment, developmental delays in children, and reduced quality of life emphasize the need for holistic, multidisciplinary care beyond vascular correction (22).

Unraveling the molecular underpinnings of MMD, such as the role of RNF213 in endothelial dysfunction or the potential of genes like ACTA2 and GUCY1A3 as therapeutic targets, is imperative for shifting from symptomatic management to disease-modifying

interventions. Emerging approaches, including gene therapy to enhance vascular stability, stem cell therapy to promote repair, and pharmacological agents like cilostazol to bolster CBF, hold promise but require rigorous clinical validation. Enhanced genetic screening, particularly in high-prevalence populations like East Asians, could refine risk stratification and personalize treatment (23). Ultimately, advancing MMD care hinges on bridging these scientific and clinical frontiers through collaborative research, innovative trials, and a deeper grasp of its physiological roots. By doing so, we can transform the prognosis of this enigmatic disorder, offering patients not just survival, but sustained neurological health and improved quality of life.

Disclosures

The authors declare no competing financial or non-financial interests in relation to the work described.

Acknowledgments

We gratefully acknowledge Dr. Gregory Shanower, Dr. Mary Pelkowski, and Dr. Michael Hardisky for their insightful discussions and valuable advice throughout this research. Their feedback significantly contributed to the clarity and direction of this work. AI was used to assist with language editing and grammar, with the final text reviewed and edited by the author and reviewers.

References

1. Cho WS, Kim JE, Kim CH, et al. Long-term outcomes after combined revascularization surgery in adult moyamoya disease. *Stroke*. 2014;45(10):3025-3031.
2. Ihara M, Yamamoto Y, Hattori Y, et al. Moyamoya disease: diagnosis and interventions. *Lancet Neurol*. 2022;21(8):747-758. doi:10.1016/S1474-4422(22)00165-X
3. Grok 3, xAI. Artificial intelligence model for data analysis and information synthesis. xAI; 2025.
4. Zou X, Liao Y, Jiang C, et al. Brain perfusion, cognition, and plasma Alzheimer's biomarkers in moyamoya disease. *Alzheimer's Dement*. 2023;19(8):3316-3326. doi:10.1002/alz.12958
5. Michiels C. Endothelial cell functions. *J Cell Physiol*. 2003;196(3):430-443. 10.1002/jcp.10333
6. Mineharu Y, Miyamoto S. RNF213 and GUCY1A3 in moyamoya disease: key regulators of metabolism, inflammation, and vascular stability. *Front Neurol*. (2021): 687088.
7. Pettersson SD, Olofsson HK, Ali S, Szarek D, Miękisiak G, Ogilvy CS. (2023). Risk factors for ischemic stroke after revascularization surgery in patients with moyamoya disease: an age-stratified comparative meta-analysis. *Wor. Neur*, 173, 146-157.
8. Kraemer M, Trakolis L, Platzen J, Schwitalla JC, Bersano A, Albrecht P, et al. Movement symptoms in European Moyamoya angiopathy—First systematic questionnaire study. *Clin Neurol Neurosurg*. 2017;152:52-56. doi: 10.1016/j.clineuro.2016.11.017
9. Funaki T, Takahashi JC, Houkin K, Kuroda S, Takeuchi S, et al. (2018). High rebleeding risk associated with choroidal collateral vessels in hemorrhagic moyamoya disease: analysis of a nonsurgical cohort in the Japan Adult Moyamoya Trial. *Jour. Neuro*, 130(2), 525-530.
10. Gao G, et al. Factors influencing collateral circulation formation after Indirect Revascularization for Moyamoya Disease: A Narrative Review. *Translational Stroke Research* 15.6 (2024): 1005-1014.
11. Kuroda S, Houkin K. Bypass Surgery for Moyamoya Disease—Concept and Essence of Surgical Techniques. *Neurologia Medico-Chirurgica* 52.5 (2012): 287-294.
12. Du L, et al. Imaging methods for surgical revascularization in patients with moyamoya disease: an updated review. *Neurosurgical Review* (2022): 1-14.
13. Yamada I, Nakagawa T, Matsushima Y, Shibuya H. High-resolution turbo magnetic resonance angiography for diagnosis of moyamoya disease. *Stroke*. 2001;32(8):1825-1829. doi: 10.1161/01.STR.32.8.1825.
14. Mertens R, et al. The genetic basis of moyamoya disease. *Translational Stroke Research* (2022): 1-21.
15. Demartini Jr. Z, et al. Moyamoya disease and

syndrome: a review. *Radiologia Brasileira* 55 (2022): 31-37.

16. Canavero I, et al. Clinical management of moyamoya patients. *Journal of Clinical Medicine* 10.16 (2021): 3628.
17. Gonzalez NR, et al. Adult moyamoya disease and syndrome: current perspectives and future directions: a scientific statement from the American Heart Association/American Stroke Association. *Stroke* 54.10 (2023): e465-e479.
18. Asselman C, et al. Moyamoya disease emerging as an immune-related angiopathy. *Trends in Molecular Medicine* 28.11 (2022): 939-950.
19. Velo M, et al. Moyamoya vasculopathy: cause, clinical manifestations, neuroradiologic features, and surgical management. *World Neurosurgery* 159 (2022): 409-425.
20. Fox BM, et al. Pathophysiology of vascular stenosis and remodeling in moyamoya disease. *Frontiers in Neurology* 12 (2021): 661578.
21. Teo M, et al. Short- and long-term outcomes of moyamoya patient post-revascularization. *Journal of Neurosurgery* 138.5 (2022): 1374-1384.
22. Gatti JR, Gonzalez Torriente A, Sun LR. Clinical presentation and stroke incidence differ by moyamoya etiology. *Journal of Child Neurology* 36.4 (2021): 272-280.
23. Singh R, et al. Research advances in the diagnosis and treatment of moyamoya disease: a bibliometric analysis. *Neurosurgical Review* 45.3 (2022): 1977-1985.